

$[\frac{1}{\sqrt{2}}(y+z)-t]$ -type and $[t\sqrt{2}/(y+z)]$ -type plane wave solutions of Einstein's field equations in V_6

^aB. G. Ambatkar , ^bJ K Jumale

^aHead Department of mathematics, R.S. Bidkar Arts, Comm. and Science College, Hinganghat, Maharashtra, India

^bR.S. Bidkar Arts, Comm. and Science College, Hinganghat, Distt. Wardha, Maharashtra, India

Abstract

Present paper deals with the study of $[\frac{1}{\sqrt{2}}(y+z)-t]$ - type and $[t\sqrt{2}/(y+z)]$ -type plane wave solutions of Einstein's field equations in general theory of relativity in the case where the zero mass scalar field coupled with gravitational field. Also zero mass scalar field coupled with gravitational and electromagnetic field and established the existence of these two types of plane wave solutions in V_6 . Furthermore, we have considered the case of massive scalar field and shown that the non-existence of these two types of plane wave solutions in GR theory.

KEYWORDS : Einstein's field equations in general relativity in case where the zero mass scalar field coupled with gravitational field and coupled with gravitational and electromagnetic field, etc.

§ 1. Introduction

H.Takeno (1961) has obtained the non-flat plane wave solutions g_{ij} of the field equations $R_{ij} = 0$ and established the existence of $(z-t)$ -type and (t/z) -type plane waves for purely gravitational case in four-dimensional empty region of space-time. On the lines of Takeno (1961), Thengane (2002) have also obtained the plane wave solutions g_{ij} of the field equations $R_{ij} = 0$ in purely gravitational case by reformulating Takeno's (1961) definition of plane wave as follows:

Definition A plane wave g_{ij} is a non-flat solution of field equations

$$R_{ij} = 0, \quad [i, j = 1,2,3,4,5,6] \tag{1.1}$$

in an empty region of space-time such that

$$g_{ij} = g_{ij}(Z), \quad Z = Z(x^i), \quad \text{where } x^i = v, u, y, z, t \tag{1.2}$$

in some suitable co-ordinate system such that

$$g^{ij}Z_{,i}Z_{,j} = 0, \quad Z_{,i} = \frac{\partial Z}{\partial x^i}, \tag{1.3}$$

$$Z = Z(y, z, t), \quad Z_{,4} \neq 0, \quad Z_{,5} \neq 0, \quad Z_{,6} \neq 0. \quad (1.4)$$

In this definition, the signature convention adopted is

$$g_{rr} < 0, \quad r = 1,2,3,4,5$$

$$\begin{vmatrix} g_{rr} & g_{rs} \\ g_{sr} & g_{ss} \end{vmatrix} > 0, \quad \begin{vmatrix} g_{rr} & g_{rs} & g_{rt} \\ g_{sr} & g_{ss} & g_{st} \\ g_{tr} & g_{ts} & g_{tt} \end{vmatrix} < 0, \quad g_{66} > 0 \quad (1.5)$$

$$[\text{not summed for } r,s=1,2,3,4,5] \text{ and accordingly } g = \det(g_{ij}) < 0 \quad (1.6)$$

In the paper [1], by investigating the plane symmetric line elements

$$ds^2 = -Adv^2 - Adu^2 - Adx^2 - B[dy^2 + dz^2 - dt^2] \quad (1.7)$$

$$\text{and } ds^2 = -Adv^2 - Adu^2 - Adx^2 - Z^2Bdy^2 - Z^2Bdz^2 + Bdt^2 \quad (1.8)$$

we have obtained the relations of non-vanishing components of Ricci tensor as

$$P = 2R_{44} = 2R_{55} = R_{66} = 2R_{45} = -\sqrt{2}R_{46} = -\sqrt{2}R_{56}, \quad (1.9)$$

$$\text{and } P = \frac{(y+z)^2}{Z^2}R_{44} = \frac{(y+z)^2}{Z^2}R_{55} = \frac{(y+z)^2}{2}R_{66} = \frac{(y+z)^2}{Z^2}R_{45}$$

$$= -\frac{(y+z)^2}{Z\sqrt{2}}R_{46} = -\frac{(y+z)^2}{Z\sqrt{2}}R_{56} \quad (1.10)$$

for $[\frac{1}{\sqrt{2}}(y+z)-t]$ -type and $[t\sqrt{2}/(y+z)]$ -type plane waves respectively.

In the present paper, we investigate whether these two types of plane wave solutions exist in the case where zero mass scalar field coupled with gravitational field and the zero mass scalar field coupled with gravitational and electromagnetic field. Furthermore we consider the coupling of massive scalar field with gravitational field and the massive scalar field with gravitational and electromagnetic field in V_6 to investigate the existence of these two types of plane wave solutions of Einstein's field equation

$$R_{ij} = (-8\pi)[T_{ij} - \frac{1}{2}g_{ij}T], \quad [i, j = 1,2,3,4,5,6] \quad (1.11)$$

where R_{ij} is the Ricci tensor of the space-time, T_{ij} is the energy momentum tensor,

$$g_{ij} \text{ is the fundamental tensor of the space-time and } T = T_i^i = g^{ij}T_{ij}.$$

We study these two plane wave solutions separately.

Case I $[\frac{1}{\sqrt{2}}(y+z)-t]$ -type type plane wave solutions in V_6

§2. Zero Mass scalar field coupled with gravitational field in V_6

The energy momentum tensor of zero mass scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k], \quad [k = 1, 2, 3, 4, 5, 6] \quad (2.1)$$

where V is scalar function of Z and $V_j = \frac{\partial V}{\partial x^j}$,

$(x^j = x^1, x^2, x^3, x^4, x^5, x^6 \text{ i.e. } x^j = v, u, y, z, t)$.

$$\text{Thus } \sqrt{2}V_4 = \sqrt{2}V_5 = -V_6 = \bar{V} \quad \because V_1 = V_2 = V_3 = 0 \quad (2.2)$$

where $(-)$ bar denotes partial derivative w. r. to Z

From line element (1.7) we have

$$g_{44} = -B, \quad g_{55} = -B, \quad (2.3)$$

$$g^{44} = -\frac{1}{B}, \quad g^{55} = -\frac{1}{B}, \quad g^{66} = \frac{1}{B} \quad (2.4)$$

$$\Rightarrow V_k V^k = 0. \quad (2.5)$$

Therefore equation (2.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j].$$

And from equation (2.6), in zero mass scalar field

$$T = T_i^i = g^{ij} T_{ij} = 0. \quad (2.7)$$

Then using equations (2.6) and (2.7), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j] \quad (2.8)$$

which further gives

$$R_{44} = R_{55} = R_{45} = -\bar{V}^2, \quad R_{66} = -2\bar{V}^2, \quad R_{46} = R_{56} = \sqrt{2} \bar{V}^2. \quad (2.9)$$

Thus non-vanishing components of Ricci tensor are related as

$$2R_{44} = 2R_{55} = R_{66} = 2R_{45} = -\sqrt{2}R_{46} = -\sqrt{2}R_{56}. \quad (2.10)$$

It is observed that the equation (2.10) is compatible with the equation (1.9) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[\frac{1}{\sqrt{2}}(y+z)-t]$ -type plane wave solutions exist in the case where zero mass scalar field is coupled with the gravitational field.

§ 3. Zero Mass scalar field coupled with gravitational & electromagnetic field in V_6

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k] + E_{ij} \quad (3.1)$$

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for zero mass scalar field

$$V_k V^k = 0 \text{ and } T = T_i^i = g^{ij} T_{ij} = 0. \quad (3.2)$$

Therefore equation (3.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j] + E_{ij}. \quad (3.3)$$

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + 4\pi E_{ij}]. \quad (3.4)$$

Then equation (3.4) yields

$$R_{44} = -2\left[\frac{\bar{V}^2}{2} + 4\pi E_{44}\right], \quad R_{55} = -2\left[\frac{\bar{V}^2}{2} + 4\pi E_{55}\right], \quad R_{66} = -2\left[\frac{\bar{V}^2}{2} + 4\pi E_{66}\right], \\ R_{45} = -2\left[\frac{\bar{V}^2}{2} + 4\pi E_{45}\right], \quad R_{46} = -2\left[-\frac{\bar{V}^2}{\sqrt{2}} + 4\pi E_{46}\right], \quad R_{56} = -2\left[-\frac{\bar{V}^2}{\sqrt{2}} + 4\pi E_{(n-1)n}\right]. \quad (3.5)$$

All above non-vanishing components of Ricci tensor are related as

$$2R_{44} = 2R_{55} = R_{66} = 2R_{45} = -\sqrt{2}R_{46} = -\sqrt{2}R_{56} \quad (3.6)$$

It is observed that the equation (3.6) is compatible with equation (1.9) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type type plane wave solutions exists in the case where zero mass scalar field is coupled with gravitational & electromagnetic field.

§ 4. Massive scalar field coupled with gravitational field in V_6

The energy momentum tensor of massive scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)], \quad [k = 1,2,3,4,5,6] \quad (4.1)$$

where V is scalar function of Z and $V_j = \frac{\partial V}{\partial x^j}$, $(x^j = v, u, x, y, z, t)$

m is mass associated with the massive scalar field.

$$\text{Thus } V_4 \sqrt{2} = V_5 \sqrt{2} = -V_6 = \bar{V} \quad \because V_1 = V_2 = V_3 = 0 \quad (4.2)$$

where (-) bar denotes partial derivative w. r. to Z .

From line element (1.7), we have obtained equations like (2.3), (2.4) and (2.5) respectively.

Therefore equation (4.1) implies

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2]. \quad (4.3)$$

Equation (4.6) yields

$$T = T_i^i = g^{ij}T_{ij} = \frac{1}{4\pi} \left[\frac{n}{2} m^2 V^2 \right]. \quad (4.4)$$

Using (4.3) and (4.4), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j - g_{ij} m^2 V^2] \quad (4.5)$$

which further yields

$$\begin{aligned} R_{44} = R_{55} &= -[\bar{V}^2 + 2Bm^2V^2], & R_{46} = R_{56} &= \bar{V}^2 \sqrt{2}, & R_{45} &= -\bar{V}^2, \\ R_{66} &= -[2\bar{V}^2 - 2Bm^2V^2]. \end{aligned} \quad (4.6)$$

But from the line element (1.7) we have the relation of non-vanishing components of Ricci tensor as

$$2R_{44} = 2R_{55} = R_{66} = 2R_{45} = -\sqrt{2}R_{46} = -\sqrt{2}R_{56}. \quad (4.7)$$

It is to be noted that here the equation (4.6) is incompatible with the equation (4.7) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[\frac{1}{\sqrt{2}}(y+z)-t]$ -type plane wave solutions doesn't exist in the case where massive scalar field is coupled with gravitational field.

Remark If $m^2 = 0$ then the equation (4.6) is compatible to (4.7) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational field.

§ 5. Massive scalar field coupled with gravitational & electromagnetic field in V_6

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)] + E_{ij} \quad (5.1)$$

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for massive scalar field

$$V_k V^k = 0 \quad \text{and} \quad T = T_i^i = g^{ij}T_{ij} = \frac{1}{4\pi} \left[\frac{n}{2} m^2 V^2 \right]. \quad (5.2)$$

Therefore equation (5.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2] + E_{ij}. \quad (5.3)$$

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j - g_{ij} m^2 V^2 + 4\pi E_{ij}]. \quad (5.4)$$

Then from equation (5.4) we have

$$\begin{aligned} R_{44} = R_{55} &= -[\bar{V}^2 + 2Bm^2V^2 + 8\pi E_{55}], & R_{66} &= -[2\bar{V}^2 - 2Bm^2V^2 + 8\pi E_{66}], \\ R_{45} &= -[\bar{V}^2 + 8\pi E_{45}], & R_{46} = R_{56} &= -[-\sqrt{2}\bar{V}^2 + 8\pi E_{46}]. \end{aligned} \quad (5.5)$$

But line element (1.7) gives the relation of non-vanishing components of Ricci tensor as

$$2R_{44} = 2R_{55} = R_{66} = 2R_{45} = -R_{46}\sqrt{2} = -R_{56}\sqrt{2}. \quad (5.6)$$

Here it has been observed that the equation (5.5) is incompatible with the equation (5.6) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[-\frac{1}{\sqrt{2}}(y+z)-t]$ -type plane wave solutions of Einstein's field equation in general relativity doesn't exist in the case where massive scalar field is coupled with gravitational and electromagnetic field.

Remark If $m^2 = 0$ then the equation (5.5) is compatible to (5.6) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational & electromagnetic field.

Case II we consider the case of $[t\sqrt{2}/(y+z)]$ -type plane wave solutions.

§ 6. Zero mass scalar field coupled with gravitational field in V_6

The energy momentum tensor of zero mass scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k], \quad [k = 1, 2, 3, 4, 5, 6] \quad (6.1)$$

where V is scalar function of Z and $V_j = \frac{\partial V}{\partial x^j}$, ($x^j = v, u, x, y, z, t$).

$$\text{Thus } V_1 = V_2 = V_3 = 0 \text{ and } V_4 = V_5 = -V_6 \frac{Z}{\sqrt{2}} = -\frac{\bar{V}Z}{(y+z)}. \quad (6.2)$$

here (-) bar denotes partial derivative w. r. to Z

From line element (1.8) we have

$$g_{44} = -Z^2 B, \quad g_{55} = -Z^2 B, \quad g_{66} = B, \quad (6.3)$$

$$g^{44} = -\frac{1}{Z^2 B}, \quad g^{55} = -\frac{1}{Z^2 B}, \quad g^{66} = \frac{1}{B} \quad (6.4)$$

$$\Rightarrow V_k V^k = 0. \quad (6.5)$$

Therefore equation (6.1)

$$\Rightarrow T_{ij} = \frac{1}{4\pi} [V_i V_j]. \quad (6.6)$$

Equation (6.6) yields

$$T = T_i^i = g^{ij} T_{ij} = 0. \quad (6.7)$$

Then from (6.6) and (6.7), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j] \quad (6.8)$$

which further gives

$$R_{44} = R_{55} = -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2}\right], \quad R_{66} = -2\left[\frac{2\bar{V}^2}{(y+z)^2}\right] \quad R_{45} = -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2}\right],$$

$$R_{46} = R_{56} = 2\sqrt{2}\left[\frac{\bar{V}^2 Z}{(y+z)^2}\right]. \tag{6.9}$$

Thus non-vanishing components of Ricci tensor are related as

$$\frac{(y+z)^2}{Z^2} R_{44} = \frac{(y+z)^2}{Z^2} R_{55} = \frac{(y+z)^2}{2} R_{66} = \frac{(y+z)^2}{Z^2} R_{45}$$

$$= -\frac{(y+z)^2}{Z\sqrt{2}} R_{46} = -\frac{(y+z)^2}{Z\sqrt{2}} R_{56}. \tag{6.10}$$

It is observed that the equation (6.10) is compatible with equation (1.10) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions exist in the case where zero mass scalar field is coupled with the gravitational field.

§ 7. Zero Mass scalar field coupled with gravitational & electromagnetic field in V_6

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi}[V_i V_j - \frac{1}{2} g_{ij} V_k V^k] + E_{ij} \tag{7.1}$$

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for zero mass scalar field

$$V_k V^k = 0 \quad \text{and} \quad T = T^i_i = g^{ij} T_{ij} = 0. \tag{7.2}$$

Therefore equation (7.1) becomes

$$T_{ij} = \frac{1}{4\pi}[V_i V_j] + E_{ij}. \tag{7.3}$$

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + 4\pi E_{ij}] \tag{7.4}$$

which further yields

$$R_{44} = -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + 4\pi E_{44}\right], \quad R_{55} = -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + 4\pi E_{55}\right],$$

$$R_{66} = -2\left[\frac{2\bar{V}^2}{(y+z)^2} + 4\pi E_{66}\right], \quad R_{45} = -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + 4\pi E_{45}\right],$$

$$R_{46} = -2\left[\frac{-\bar{V}^2 Z\sqrt{2}}{(y+z)^2} + 4\pi E_{46}\right], \quad R_{56} = -2\left[\frac{-\bar{V}^2 Z\sqrt{2}}{(y+z)^2} + 4\pi E_{56}\right]. \tag{7.5}$$

All above non-vanishing components of Ricci tensor are related as

$$\begin{aligned} \frac{(y+z)^2}{Z^2} R_{44} &= \frac{(y+z)^2}{Z^2} R_{55} = \frac{(y+z)^2}{2} R_{66} = \frac{(y+z)^2}{Z^2} R_{45} \\ &= -\frac{(y+z)^2}{Z\sqrt{2}} R_{46} = -\frac{(y+z)^2}{Z\sqrt{2}} R_{56}. \end{aligned} \quad (7.6)$$

It is observed that the equation (7.6) is compatible with equation (1.10) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions exist in the case where zero mass scalar field is coupled with gravitational & electromagnetic field.

§ 8. Massive scalar field coupled with gravitational field in V_6

The energy momentum tensor of massive scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)], \quad [k = 1, 2, 3, 4, 5, 6] \quad (8.1)$$

where V is scalar function of Z and $V_j = \frac{\partial V}{\partial x^j}$, ($x^j = v, u, x, y, z, t$),

m is mass associated with the massive scalar field.

$$\text{Thus } V_4 = V_5 = -V_6 \frac{Z}{\sqrt{2}} = -\frac{\bar{V}Z}{(y+z)}, \quad \therefore V_1 = V_2 = V_3 = 0 \quad (8.2)$$

where (-) bar denotes partial derivative w. r. to Z .

From line element (1.8), we have obtained equations like (6.3), (6.4) and (6.5).

Therefore equation (8.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2]. \quad (8.3)$$

From equation (8.6) we have

$$T = T^i_i = g^{ij} T_{ij} = \frac{1}{4\pi} [3m^2 V^2]. \quad (8.4)$$

Then from (8.3) and (8.4), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j - g_{ij} m^2 V^2] \quad (8.5)$$

which further yields

$$\begin{aligned} R_{44} = R_{55} &= -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + Z^2 B m^2 V^2\right], & R_{45} &= -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2}\right], \\ R_{66} &= -2\left[\frac{2\bar{V}^2}{(y+z)^2} - B m^2 V^2\right], & R_{46} = R_{56} &= -2\left[\frac{-\bar{V}^2 Z\sqrt{2}}{(y+z)^2}\right]. \end{aligned} \quad (8.6)$$

But from the line element (1.8) we have the relation of non-vanishing components of Ricci tensor as

$$\begin{aligned} \frac{(y+z)^2}{Z^2} R_{44} &= \frac{(y+z)^2}{Z^2} R_{55} = \frac{(y+z)^2}{2} R_{66} = \frac{(y+z)^2}{Z^2} R_{45} \\ &= \frac{-(y+z)^2}{Z\sqrt{2}} R_{46} = \frac{-(y+z)^2}{Z\sqrt{2}} R_{56} \end{aligned} \quad (8.7)$$

Here we observed that the equation (8.6) is incompatible with the equation (8.7) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions of Einstein's field equation in general theory of relativity doesn't exist in the case where massive scalar field is coupled with the gravitational field.

Remark If $m^2 = 0$ then the equation (8.6) is compatible to (8.7) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational field.

§ 9. Massive scalar field coupled with gravitational & electromagnetic field in V_6

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi}[V_i V_j - \frac{1}{2} g_{ij}(V_k V^k - m^2 V^2)] + E_{ij} \quad (9.1)$$

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for massive scalar field

$$V_k V^k = 0 \quad \text{and} \quad T = T^i_i = g^{ij} T_{ij} = \frac{1}{4\pi} [\frac{n}{2} m^2 V^2]. \quad (9.2)$$

Therefore equation (9.1) becomes

$$T_{ij} = \frac{1}{4\pi}[V_i V_j + \frac{1}{2} g_{ij} m^2 V^2] + E_{ij}. \quad (9.3)$$

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j - g_{ij} m^2 V^2 + 4\pi E_{ij}]. \quad (9.4)$$

And from equation (9.4) we have

$$\begin{aligned} R_{44} &= -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + Z^2 B m^2 V^2 + 4\pi E_{44}\right], & R_{55} &= -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + Z^2 B m^2 V^2 + 4\pi E_{55}\right], \\ R_{66} &= -2\left[\frac{2\bar{V}^2}{(y+z)^2} - B m^2 V^2 + 4\pi E_{66}\right], & R_{45} &= -2\left[\frac{\bar{V}^2 Z^2}{(y+z)^2} + 4\pi E_{45}\right], \\ R_{46} &= -2\left[\frac{-\bar{V}^2 Z \sqrt{2}}{(y+z)^2} + 4\pi E_{46}\right], & R_{56} &= -2\left[\frac{-\bar{V}^2 Z \sqrt{2}}{(y+z)^2} + 4\pi E_{56}\right]. \end{aligned} \quad (9.5)$$

But line element (1.8) gives the relation of non-vanishing components of Ricci tensor as

$$\begin{aligned} \frac{(y+z)^2}{Z^2} R_{44} &= \frac{(y+z)^2}{Z^2} R_{55} = \frac{(y+z)^2}{2} R_{66} = \frac{(y+z)^2}{Z^2} R_{45} \\ &= \frac{-(y+z)^2}{Z\sqrt{2}} R_{416} = \frac{-(y+z)^2}{Z\sqrt{2}} R_{56}. \end{aligned} \quad (9.6)$$

Here we observed that the equation (9.5) is incompatible with the equation (9.6) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions of Einstein's field equation doesn't exist in the case where massive scalar field is coupled with gravitational and electromagnetic field.

Remark If $m^2 = 0$ then the equation (9.5) is compatible to (9.6) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational and electromagnetic field.

References

- [1] B. G. Ambatkar : “ $[\frac{1}{\sqrt{2}}(y+z)-t]$ -type and $[t\sqrt{2}/(y+z)]$ -type four and J K Jumale (2015) dimensional plane wave solutions of Einstein's field equations in general relativity”. *Research Inventory Vol. 5, Issue 2, PP-30-38.*
- [2] I.S.Mohurleya, R. K. Jumale : “ $[\frac{1}{\sqrt{2}}(y+z)-t]$ -type and $[t\sqrt{2}/(y+z)]$ -type five and J K Jumale (2014) dimensional plane wave solutions of Einstein's field equations in general relativity”. *Online International Interdisciplinary Research Journal, Vol.-IV, Issue-IV.*
- [3] H Takeno(1961) : “The Mathematical theory of plane gravitational wave in general relativity”. *Scientific report of the Research Institute to theoretical physics. Hiroshima, University, Takchara, Hiroshima-ken, Japan-1961.*