

Mimusops elengi* Linn.: A Potential Phytochemical Agent in Clinical Periodontics*Akanksha choudhary^a, Smitha C N^b, Suresh D K^c, Ajay Kumar Srivastava^d**^aPG Student; ^bProfessor; ^cProfessor and Head; Department of Periodontology and oral Implantology, ; Maharishi Markandeshwar College of Dental Sciences and Research, Mullana, Ambala, Haryana, India^dHead, Dept. of Botany, St. Xavier's College, Ranchi, Jharkhand, India**Abstract**

Ayurvedic medicine is an integral part of Rigveda. More than 80% of population in the developing countries use traditional or local medicine, which is originally a part of ayurveda. Natural medicines are considered to be “safe”, as they are natural, without any synthetic chemical. These herbs are very much popular because of their less toxicity and adverse effects. One of these herbs, a common Indian native plant is *Mimusops elengi* Linn. For a long time in the traditional ayurveda system of medicine, *Mimusops* has been described and used for its potential medicinal values. Various parts of this plant such as leaves, twig, bark, stem, root, fruits, seeds, and flowers have shown improvement in oral health when used in different concentrations or in combination with other plant materials. Ethanolic extract of bark shows anti-inflammatory, analgesic as well as antipyretic effect. The phytochemical agents such as Lupeol, saponin present in the bark show anti-inflammatory activity and tannins show antibacterial property against different strains of *Streptococcus* species, *Staphylococcus* species and *E. fecalis*. The extract of bark in different materials has shown significant antioxidant potential and scavenging property. This plant should be explored and further research work should be carried out in periodontal field for new valuable drug formulation, which will be economically more suitable to the majority of populations in developing countries.

KEYWORDS: drug formulation, local medicine, *Mimusops elengi*, phytochemical agents.

Introduction:

Ayurvedic medicine is an integral part or supplement of Rigveda. It is considered as Upaveda i.e. auxiliary knowledge in the ancient vedic system. The main text of ayurvedic medicine are Charakasamhita and Sushrutasamhita, which were written during 100 BC and the field was flourished throughout the Indian subcontinent, including present countries like India, Pakistan, Afghanistan, Srilanka, Iraq, Iran. According to WHO survey, upto 80% of population in the developing countries use some form of traditional or local medicine, which is originally a part of ayurveda.¹ The central council of Indian medicine (CCIM), 1971 has established The Department of Ayurveda, Yoga and Naturopathy, Unani, Siddha and Homeopathy (AYUSH) as an integral part of Ministry of health and family welfare, Government of India.² Ayurvedic tradition is mainly based on the use of plant medicines. Modern medicinal system in all over the world has recognized the importance of herbal medicine, also known as herbalism or herbology. Natural medicines are considered to be “safe”, as they donot contain any synthetic chemicals. These herbs are very much popular as they are easily available with less toxicity and adverse effects.

One of these herbs, a common Indian native plant is *Mimusops elengi* Linn. It is a handsome evergreen tree of family Sapotaceae; Linn stands for “Carl Linnaeus” who first described the modern taxonomy of plants. It is also called as Bulletwood tree or Spanish cherry in English; Bakul in Hindi; Maulsari or Chirapushpa in Sanskrit; Elengi in Malayalam; Ranja in Kanada.³ It reaches upto a height of 15-16 meter, found in all over the Indian subcontinent.⁴ The new leaves come generally in the month of February and shape is oval, elliptical, with undulated margin. The leaves are 6-10 cm. long and 3-5 cm. wide with vibrant shiny green coloured surface. The older leaves become dark green in colour. The tree has grayish black channeled bark, which has 10-20 cm longitudinal fissures. The tree is sacred in India since ancient time, due to its fragrant flowers, which are widely used in gardens, temples, ornaments of Hindu Gods, garlands etc. These white flowers have been described as symbol of love and beauty in famous Sanskrit literature of Kalidasa.⁵ The berry is yellow-orange in colour, fruiting in august and is having astringent property. Seeds are grayish-brown in colour.



Fig.1: Stem bark

PHYTOCHEMICAL EXTRACT OF MIMUSOPS ELENGI LINN.

Stem bark:

The bark contains alkaloids, starch, saponins, tannin, some caoutchoue, wax coloring matter, starch and ash forming inorganic salts.⁶ The major constituents of bark are Taraxerone, taraxerol, betulinic acid spinasterol, sodium salt of betulinic acid and ursolic acid, Fatty acid esters of alpha-spinasterol.⁷ Along with the triterpenoids, farnan-3-one, and olean-18-en-2-one-3-ol and lup-20 (29)-en-3 beta-ol, mimusopfarnanol, a new chemical i.e. farnanetypepentacyclic triterpene, farnan-2-one-3 betaol, has also been isolated.⁸ Another study showed the isolation of beta amyryn, lupeol (triterpene 3 β -hydroxy-lup-20(29)-ene-23, 28-dioic acid).⁹ Ruikar et al(2009)¹⁰ studied and collected sample of 0.18% volatile organic matter from the steam distillation of bark. Gallic acid esters, known as phenyl propyl gallate, alpha cadinol, taumurolol, hexadecanoic acid, diisobutyl phthalate, octadecadienoic acid are also reported to be obtained from the bark of *Mimusops*.¹¹

Fruits and seeds:

Contents of fruits are moisture (79.27 %), protein (1.29%), fat (2.76 K Cal), sugar (15.2%), Fiber (1.13%), Vitamin C (3.27 mg / 100 gm), Mineral content (0.32%), Iron


(0.59 mg / 100 gm) , Sodium (5.16 mg / 100 gm) , Potassium (98.54 mg / 100 gm).¹² Different types of acids isolated from fruits are Quercitol, ursolic acid, dihydroquercetin, quercetin, β - d glycosides of β sitosterol, alphaspinasterol.⁷ Two new Pentacyclic triterpene acids have also been isolated which are named asmimusops acid and mimusopic acid. These two acids contain oleanane skeleton and mimusopane.¹³ These acids contain the mimusops gene and mimugene.¹⁴


The seeds of Mimusops contain novel triterpenoid saponins, named as mimusops in and saponinmimus in,¹⁵ triterpenoid saponins, Mi-saponin A and 16 α -hydroxy Mi-saponin A.¹⁶ some new types of saponins are also isolated which are taxifolin, α -spinasterol glucoside, Miglycoside-1, triterpenoid saponins, mimusopside A and B.^{17,18}

Leaves and roots:

The ethanolic extract of Mimusops leaves has been isolated and its chemical contents are quercitol, hentriacontane, β -carotene, glucose, D-mannitol, β -sitosterol, β -sitosterol- β -D-glucoside.^{19,20,21} The roots of Mimusops contain steroidal saponins 5 α -stigmast-9(11)en-3-o-beta-D-glucopyranosyl (1-5)-o-beta-D-xylofuranoside.^{19,22}

CHEMICAL STRUCTURE OF MAJOR CONSTITUENTS OF MIMUSOPS ELENGI LINN.²³





SAPONIN²⁴

POTENTIAL EFFECTS OF PLANT ON VARIOUS DENTAL DISEASES:

For a long time in the traditional ayurveda system of medicine, Mimusops has been described and used for its potential medicinal values. All the different parts of this plant such as leaves, twig, bark, stem, root, fruits, seeds, and flowers are used in different concentration or in combination with other plant materials to improve oral health.

Since ancient time, Mimusops has been used as an astringent to strengthen the teeth and gingiva. The root, bark and twig are used as “*dantwan*” to clean the teeth and tongue in rural areas of India.²⁵ The stem or bark has been found to be the most effective and potent part showing astringent, antimicrobial, antioxidant and anti-inflammatory property. The bark mixed with pepper, honey and ghee (*kawath*) is used as a gargle to reduce gingival inflammation and strengthen the teeth. The combination of root bark decoction taken along with milk for three days in the morning is said to strengthen the teeth in older individuals.²⁶ One of the ayurvedic herbal preparation named as “Mahakhadiradivati”, which is indicated for swollen and inflamed gingiva, bad odour, stomatitis, contains Mimusops bark as an active ingredient.²⁶

Ethanol extract of bark shows anti-inflammatory, analgesic as well as antipyretic effect.²⁷ The phytochemical agents such as Lupeol, saponin present in the bark show anti-inflammatory activity and tannins show antibacterial as well as antioxidant properties.²⁸ The extract of bark in different materials has shown significant antioxidant potential and scavenging property. The ethanolic, methanolic, acetone, petroleum and water extract of bark showed antibacterial property against different strains of Streptococcus species, Staphylococcus species and against *E. fecalis*.^{29,30} The fruit and seed extracts are found to be less active, in comparison to bark extract against gingival diseases and oral pathogens.³¹

It has been used as an active ingredient in a variety of herbal toothpastes like, Vicovajradanti®, Sudantha® and Aloe dental cream® for their antimicrobial and astringent properties. The oral hygiene and gingival health have been found to be improved, as well as different clinical parameters such as bleeding on probing, plaque index and probing pocket depth have also been reduced by the use of the herbal toothpaste containing Mimusops.³²

In a study by Mistry et al (2014)³³, different concentration of Mimusops bark extract has been examined against *E. fecalis*, *Streptococcus mutans* and *Staphylococcus aureus*. 3 mg concentration was found to produce the maximum inhibitory concentration (MIC) against *E. fecalis*.

Prabhat et al in 2010³⁴ compared the petroleum ether, acetone, methanol and water extracts of bark, which were tested for their antimicrobial effect against five oral microorganisms *Staphylococcus aureus*, *Streptococcus mutans*, *Streptococcus sanguis*, *Streptococcus salivarius*, *Lactobacillus acidophilus* and *Candida albicans*. Methenolic and water extract were found to be more effective against all strains. The Acetone and ethenol extracts of bark were tested in children for their antibacterial effect against salivary microflora by “paper disc diffusion method”. The extracts were found to be effective against different *staphylococcus* species.³⁵

Different extracts of bark such as petroleum ether, benzene, chloroform, acetone, methanol and water by saxchlation process have been prepared and were examined against gram positive and gram negative bacteria of dental plaque and calculus by ditch plate method. The best result was obtained by chloroform extract.³⁶ Kulkarni AA et al³⁷ in their study stated, that aqueous and acetone extracts of bark act against salivary microflora, which was evaluated by paper disc diffusion method. The results were not significantly different for both the extracts, although extracts were active against the pathogens.

Malhotra et al in 2011³⁸ compared the efficacy of a commercially available herbal mouthrinse with chlorhexidine gluconate, in which *Mimusops* bark was used for its astringent property. Though the herbal mouthrinse was less effective than chlorhexidine gluconate, but the clinical parameters such as bleeding on probing and plaque index showed significant difference from baseline. Swelling and inflammation of gingiva have been reduced clinically.

The ethanolic extract and β amyriacaprylate was compared with Indomethacin and the extract was found to show anti-inflammatory property.³⁹ Phogat et al in 2014²⁸ compared the Chlosite gel and herbal gel with SRP alone. Results indicated significant reduction in all the clinical parameters, such as plaque index, gingival index, probing pocket depth and clinical attachment levels at the sites where the local application of herbal gel alongwith SRP was done. The herbal gel was containing *Mimusops* as an active anti-inflammatory, antibacterial and anti-oxidant ingredient.

The chloroform and methanol extracts of bark were analysed by 1,1-diphenyl-2-picrylhydrazyl or DPPH radical, nitric oxide and some other hydroxyl radicals and the results showed that the extracts had significant antioxidant potential and scavenging property.^{40,41}

In a different in-vivo study, the petroleum, chloroform and alcohol extracts were compared for their antioxidant property against lipid peroxidation, superoxide dismutase and catalase activity. It has been found that alcohol extract was the most potent antioxidant agent in comparison to other two extracts.⁴²

CONCLUSION:

M. elengi Linn. has shown so many medicinal as well as periodontal properties. It has also been found to be more effective against various species of oral pathogens. Despite these studies and its easy availability in the society, it has not been extensively used in periodontal field. All the parts of the plant have been found useful with their medicinal value, but this plant should be explored more and further research work should be carried out in dental and periodontal field for new valuable and effective drug formulation, which will be economically and easily more suitable to the majority of populations in southern Asiatic countries.

References:

1. Legal status of traditional medicine and complementary/alternative medicine: a worldwide review. World health organization (WHO) source; 2014.
2. Welcome to central council for research in ayurveda and siddha (CCRAS); 2012.
3. Reddy MB, Reddy KR, Reddy MN. A survey of plant crude drugs of Anantapur district, Andhra Pradesh, India. *Int J Crude Drug Res.* 1989; 27(3): 141-155.
4. Kritkar KR, Basu BD. *Indian medicinal plants.* 2001;2nd Ed. Vol. II: 1224-1227.
5. Mitra R. Bakula- A reputed drugs of ayurveda, its Hist.-uses in Indian medicine. *Ind J HistSci* 1981; 16(2):169-180.
6. Gami B, Parabia MH. Pharmacognostic evaluation of bark and seeds of *Mimusopselengi*Linn. *Int J Pharm Pharmaceut Sci.* 2010; 2(4): 110-113.
7. Mishra G, Mitra CR. Constituents of bark of *Mimusopselengilinn.* *Phytochem* 1967; 6: 1909.
8. Akhtar N, Ali M, Alam MS. Pentacyclic triterpenes from the stem bark of *Mimusopselengilinn.* *Acta Pol Pharm* 2009; 66(5):549-552.
9. Jahann N, Malik A, Mustafa G, Ahmad Z, Ahmad S, Anis E *et al.* Triterpenes From *Mimusopselengi.* *Nat Prod Lett* 2001; 15(3):177-185.
10. Ruikar, Torane R, Tambe A, Puranik V, Deshpande N. GC-MS study of a steam volatile matter from *Mimusopselengi.* *Int J Chemtech Res Coden* 2009; 1(2):158-161.
11. Akhtar N, Ali M, Alam MS. Gallic acid esters from the stem bark of *Mimusopselengil.* *Nat Prod Res* 2010; 24(10):962-72.
12. Nazarudeen A. Nutritional composition of some lesser-known fruits used by the ethnic communities and local folks of kerala. *Ind J Traditional Knowledge* 2010; 9(2):398-402.
13. Sen S, Sahu NP, Mahato SB. Novel migrated oleananetriterpenoidsapogenins from *Mimusopselengi.* *Tetrahedron* 1993; 49(40):9031-9038.
14. Sen S, Sahu NP, Mahato SB. Pentacyclictriterpenoids from *Mimusopselengi.* *Phytochem* 1967; 38(1):205-207.
15. Sahu NP, Koike K, Jia Z, Nikaido T. Novel triterpenoidsaponins from *Mimusopselengi.* *Tetrahedron* 1995; 51(48):13435-13446.
16. Sahu NP, Koike K, Jia Z, Nikaido T. Triterpenoidsaponins from *Mimusopselengi.* *Phytochem* 1997; 44(6):1145-1149.
17. Sahu NP. Triterpenoidsaponins of *Mimusopselengi.* *Phytochem* 1996; 41(3):883-886.
18. Lavaud C, Massiot G, Becchi M, Misra G, Nigam SK. Saponins from three species of mimusops. *Phytochem* 1996; 41(3):887-893.
19. Manjeshwar SB, Ramakrishna JP, Harshit PB, Princy LP, Rekha B. Chemistry and medicinal properties of Bakul (*Mimusopselengi*Linn). *Rev Food Res Int.* 2011; 44: 1823-1829.
20. Kalitha D, Saikia CN. Chemical constitute and energy content of same latex bearing plants. *Biores Technol.* 2004; 92(3): 219-227.
21. Gami B. 2007. Evolution of pharmacognosticantihemorrhoidal properties of *Mimusopselengi*Linn, Ph. D. thesis Veer Narmad South Gujrat University, India.

22. Saxena VK, Srivastava K. New steroidal saponins from the roots of *Mimusopselengi*. *Fitoterapia*. 1988; 59(5): 418.
23. Kadam PV, Yadav KN, Deoda RS, Shivatare RS, Patil MJ. *Mimusopselengi*: a review on ethanobotany, phytochemical and pharmacological profile. *J PharmacognPhytochem*. 2012; 1(3): 71-81.
24. Singh KL, Srivastava P, Kumar S, Singh DK, Singh VK. *Mimusopselengi* Linn. (Maulsari): a potential medicinal plant *Archives of Biomedical Sciences* 2014; 2 (1): 18-29.
25. Kalita D, Saikia CN. Chemical constitute and energy content of some latex bearing plants. *BioresourceTechnol*2004; 92(3): 219-227.
26. Gami B, Pathak S, Parabia M. Ethnobotanical, phytochemical and pharmacological review of *Mimusopselengi*Linn. *Asian Pac J Trop Biomed* 2012; 2(9): 743-748.
27. Purnima BC, Koti AH, Thippeswamy MS, Jaji AH, Swamy YV, Kurhe AJ *et al*. Antiinflammatory, analgesic and antipyretic activities of *Mimusopselengilinn*. *Ind j pharm sci* 2010; 72(4):480-485.
28. Phogat M, Rana T, Prasad N, Baiju CS. Comparative evaluation of subgingivally delivered xanthan-based chlorhexidine gel and herbal extract gel in the treatment of chronic periodontitis. *J Indian SocPeriodontol* 2014;18:172-7.
29. Jebashree HS, Kingsley SJ, Sathish ES, Devapriya D. Antimicrobial Activity of Few Medicinal Plants against Clinically Isolated Human Cariogenic Pathogens—An *In vitro* Study. *ISRN Dentistry* 2011; 67-72.
30. Mistry KS, Sanghvi Z, Parmar G, Shah S The antimicrobial activity of *Azadirachtaindica*, *Mimusopselengi*, *Tinosporacardifolia*, *Ocimum sanctum* and 2% chlorhexidine gluconate on common endodontic pathogens: An *in vitro* study *European Journal of Dentistry*, Vol 8 / Issue 2 / Apr-Jun 2014.
31. Shahwar D, Raza MA. *In vitro* antibacterial activity of extracts of *Mimusopselengi* against gram positive and gram negative bacteria. *Afr J Microbiology Res* 2009; 3(8):458-462.
32. Jayashankar S, Panagoda GJ, Amaratunga E, Perera K, Rajapakse PS. A randomised double-blind placebo-controlled study on the effects of a herbal toothpaste on gingival bleeding, oral hygiene and microbial variables. *Ceylon Medical Journal* 2011;56(1):5-9.
33. Gupta N, Jain UK. Investigation of wound healing activity of methanolic extract of stem bark of *Mimusopselengilinn*. *Afr J Trad, Compl& Alt Med*2011; 8(2): 98-103.
34. Prabhat, Ajaybhan, Navneet, Chauhan A. Evaluation of Antimicrobial Activity of Six Medicinal Plants against Dental Pathogens. *Report Opinion* 2010; 2(6):37-42.
35. Deshpande RR, Ruikar A, Panvalkar PS, Kulkarni AA, Khatiwora E, Adasul V *et al*. Comparative evaluation of different concentrations of *Mimusopselengi*(L) extract as an antimicrobial agent against salivary micro flora. *J Biomed Sci and Res* 2010; 2(3):151-154.
36. Murudkar SS, Mundhada PA. Tatke. Antibacterial activity of *Mimusopselengilinn* bark against dental pathogens. *Ind J Pharm Educ Res* 2007; 41 (2):114-120.
37. Kulkarni AA, Deshpande RR, Panvalkar P, Mahajan P, Kale A, Ruikar A *et al*. Comparative evaluation of antibacterial properties of different extracts of

- Mimusopselengi*(bakul) & *juglansregia*(walnut) against salivary microflora. Res J PharmaceutBiolChemSci 2011; 2(3):635.
38. Malhotra R, Grover V, Kapoor A, Saxena D. Comparison of the effectiveness of a commercially available herbal mouthrinse with chlorhexidine gluconate at the clinical and patient level. Journal of Indian Society of Periodontology 2011;15(4).
 39. Rajkumara S, Pandiselvi A, Sandhiya G. Isolation of Chemical Constituents from *Mimusopselengi* bark and Evaluation of Anti-inflammatory activity. Int J Phytopharmacy 2012; 3(1):9-15.
 40. Rao KS, Munjuluri PR, Keshar NK. *In vitro* antioxidant activity and total phenolic content of *Mimusopselengi* bark. Ind J Pharm Edu Res. 2011; 45(4):317-323.
 41. Shaik J, Khasim SM, Naidu PB. Protective activity of ethanolic leaf extract of *Mimusopselengilinn* on lipid peroxidation and antioxidant enzymes in experimental diabetic rats. PHARMANEST – Int J Advances PharmaceutSci 2011; 2(2):264-275.
 42. Ashok P, Koti BC, Vishwanathswamy AH. Antirolithiatic and antioxidant activity of *Mimusopselengi* on ethylene glycol-induced urolithiasis in rats. Ind J Pharmacol 2010; 42(6): 380-383.